



Protocol 2000

Protocol 2000 is, in essence, our Cancer Protocol, but we are not naming it "Cancer Protocol" as such because it also works well for most other life-threatening diseases. I

have observed that more than 90% of those who use Protocol 2000 faithfully, and take responsibility for using it as directed here, overcome their cancer or other disease completely. However, I must also mention that there are cases of cancer and other diseases that simply are too far gone for even MMS1 and MMS2 to help. Normally these are the cases that have had tremendous amounts of chemo, radiation, or surgery treatment and the body is simply “past the point of no return.” However, we never say never. If the person still has one more hour to live, get some MMS1 into him. See page 249 for more information on helping people with extreme conditions.

On Protocol 2000 you will:

- Increase the number of drops you take each hour to as many drops as you can handle (up to the maximum amount of drops for your weight—see page 92) without getting sick due to the MMS. In most cases the increase in drops is needed for cancer and other life-threatening diseases.
- Increase the number of hours you take your dose each day from eight to ten hours.
- At the beginning of the third or fourth day of Protocol 2000, you should begin taking MMS2 in addition to MMS1.

The most important thing to remember is, **never stop taking MMS until you are well.** Remain on Protocol 2000 and any needed Supporting Protocols, as explained in the Health Recovery Plan, until you have fully recovered your health.

Note: *When progressing from Protocol 1000 Plus to Protocol 2000, always continue using DMSO according to the instructions given in Protocol 1000 Plus while doing*

Protocol 2000, i.e. add 3 drops of DMSO to the dose for every 1 drop of MMS1.

Instructions for Protocol 2000

Step 1

- ❑ Increase the number of hours per day that one takes the hourly dose to ten hours per day instead of the eight hours per day of Protocol 1000.

Step 2

- ❑ Begin increasing the drops in your daily dose by 1 drop increments. For example, if you were taking 3 drops an hour as per Protocol 1000, you can increase to 4 drops.
- ❑ The Health Recovery Plan (HRP) gives allowance for an exception to the rule, (see page 70). If you fall into this category and therefore come to Protocol 2000 directly from the Starting Procedure because of cancer or some other life-threatening disease, then begin at 1 drop per hour and increase the drops per hour after only a few hours at 1 drop per hour. You can tell if you should not add another drop per hour by the way you feel. Just keep increasing by 1 additional drop per hour until a tiny sickish feeling beyond how the disease makes you feel, lets you know for the time being to stop increasing. Some people can move along quicker and some cannot, please be attentive to the Three Golden Rules of MMS.
- ❑ It is important to not allow yourself to feel worse than your disease is already making you feel, as the additional sickness can then slow your recovery down. So if taking your MMS dose results in nausea, vomiting, diarrhea or excessive tiredness reduce the number of

drops you are taking by 50% for the next dose, if it still seems like the MMS1 is continuing to cause distress, then decrease the dose by another 50% of what you are taking. When you feel comfortable with the amount of MMS1 you are taking, then slowly increase the drops again. If the added sickness is severe then temporarily stop taking the drops altogether and start again as soon as you are feeling better. And again, increase to as much as you can take without feeling worse than you already are.

The following chart gives the theoretical maximum amount of drops that most people should take for their body weight. Anyone weighing more than 200 pounds can calculate their maximum number of drops by adding 1 drop for each 20 pounds over 200 pounds. There are times when a cancer is not improving that one might go ahead and take more drops per hour than suggested here, in that case do not hesitate to do so, but normally this chart is correct. Remember, follow the Three Golden Rules of MMS. Some people will not get up to anywhere near these amounts. These are maximum amounts—they are not a goal.

Protocol 2000 —Maximum MMS1 Dosage	
Weight	Dosage
80-100 lbs (36-45 kg)	Take no more than 8 drops hourly
100-120 lbs (45-54 kg)	Take no more than 8 drops hourly
120-140 lbs (54-63 kg)	Take no more than 9 drops hourly
140-160 lbs (63-72 kg)	Take no more than 10 drops hourly
160-180 lbs (72-81 kg)	Take no more than 11 drops hourly
180-200 lbs (81-90 kg)	Take no more than 12 drops hourly
200 lbs (90 kg) and above: Increase the maximum dose by 1 drop for each additional 20 lbs (9 kg)	

Step 3

- ❑ Begin taking MMS2 on the third or fourth day into the protocol. (Please read section MMS2—Details, page 274, for information on where to purchase calcium hypochlorite, and instructions on how to make MMS2 capsules.)
- ❑ Use either #1 size capsules which are the smallest that you should use, or #0 size capsules, which is one size larger than #1. (And no, I didn't make a mistake on capsule sizes; they really get smaller in size as you increase the number.) Start by loading the #1 size capsules 1/8 full or #0 size capsule about 1/16 full. When the capsules are pulled apart, one side is always larger than the other side. Fill the larger side (pack loosely). Then put the smaller side on and be sure you push it down securely in place.

Step 4

- ❑ Step 3 gets you started, but increase the amount you put in the capsule **over the next several days**, working up to either full for #1 size capsule, or 3/4 full for a #0 size capsule. Slowly increase the amount you put in the capsule.

Step 5

- ❑ Take one of these capsules 5 times a day—once every two hours.
- ❑ Take your first MMS2 capsule one-half hour after taking your second MMS1 dose of the day.

Sample Time Schedule for Protocol 2000, Once MMS2 is Added to Your Dosing

Protocol 2000 MMS2 Time Schedule	
Time	Dose
9:00 AM	MMS1 dose
10:00 AM	MMS1 dose
10:30 AM	MMS2 dose
11:00 AM	MMS1 dose
12:00 PM	MMS1 dose
12:30 PM	MMS2 dose
1:00 PM	MMS1 dose
2:00 PM	MMS1 dose
2:30 PM	MMS2 dose
3:00 PM	MMS1 dose
4:00 PM	MMS1 dose
4:30 PM	MMS2 dose
5:00 PM	MMS1 dose
6:00 PM	MMS1 dose
6:30 PM	MMS2 dose

Notes

➤ *While working up to the correct size capsules of MMS2, which is either a full #1 size capsule, or a 3/4 full #0 size capsule (**never go beyond these amounts**), keep your MMS1 doses constant. In other words, do not be working on increasing your drops of MMS1, while you are working up to your proper dose of MMS2, because if you get nauseous you will not be able to determine which of the two might be causing you to feel sick. Once you have reached the suggested amount of MMS2, then you can begin increasing your drops of MMS1 once more.*

- *Remember, at any time, whether you are increasing your amounts of MMS1, or MMS2, if at any time you feel nauseous or sick from the increase, decrease the amount by at least one half and build back up slowly.*

- **Never take a dose containing DMSO and an MMS2 capsule at the same time!** See pages 23-24 for the full warning on this.